



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports of States and yearly and monthly reports of cities.

CALIFORNIA—*Sacramento*.—Month of March, 1897. Estimated population, 30,000. Total deaths, 44, including phthisis pulmonalis, 6; diphtheria, 1, and measles, 1.

FLORIDA—*Pensacola*.—Month of March, 1897. Estimated population, 15,000. Total deaths, 20, including 1 from enteric fever.

Tampa.—Month of March, 1897. Estimated population, 26,000. Total deaths, 25, including 4 from phthisis pulmonalis.

INDIANA—*Evansville*.—Month of January, 1897. Estimated population, 65,000. Total deaths, 79, including phthisis pulmonalis, 8; enteric fever, 1; diphtheria 2, and measles, 2.

Month of February, 1897. Total deaths, 67, including phthisis pulmonalis, 6; measles 2, and whooping cough, 3.

Month of March, 1897. Total deaths, 62, including phthisis pulmonalis, 6; enteric fever, 1, and diphtheria, 1.

KENTUCKY—*Louisville*.—Month of March, 1897. Estimated population, 215,572. Total deaths, 329, including phthisis pulmonalis, 62; enteric fever, 5; diphtheria, 1; croup, 2, and whooping cough, 2.

LOUISIANA—*Shreveport*.—Month of March, 1897. Estimated population—white, 7,500; colored, 7,500; total, 15,000. Deaths—white, 4; colored, 11, including phthisis pulmonalis, 3, and enteric fever, 1.

MASSACHUSETTS—*Newton*.—Month of March, 1897. Estimated population, 27,590. Total deaths, 36, including phthisis pulmonalis, 3, and diphtheria, 1.

MICHIGAN.—Week ended April 3, 1897. Reports to the State board of health, Lansing, from 59 observers indicate that intermittent fever and erysipelas increased in area of prevalence. Phthisis pulmonalis was reported present during the week at 161 places, measles at 76, diphtheria at 30, scarlet fever at 5, enteric fever at 13, whooping cough at 7, and smallpox at 1 place (Blissfield Township).

NEW HAMPSHIRE.—*Concord*.—Month of March, 1897. Estimated population, 19,000. Total deaths, 47, including 5 from phthisis pulmonalis.

OHIO—*Toledo*.—Month of March, 1897. Estimated population, 137,780. Total deaths, 152, including phthisis pulmonalis, 30; enteric fever, 2; diphtheria, 1; measles, 4; croup, 3, and whooping cough, 2.

RHODE ISLAND—*Newport*.—Month of March, 1897. Estimated population, 21,500. Total deaths, 39, including phthisis pulmonalis, 3; croup, 1; enteric fever, 1, and whooping cough, 2.

TENNESSEE—*Chattanooga*.—Month of March, 1897. Estimated population—white, 27,000; colored, 13,000; total, 40,000. Deaths—white, 18; colored, 24; total, 42, including phthisis pulmonalis, 4, and enteric fever, 2.

Memphis.—Month of March, 1897. Estimated population—white,

28,573; colored, 27,561; total, 56,134. Deaths—white, 37; colored, 55; total, 92, including 16 from phthisis pulmonalis.

WEST VIRGINIA—*Wheeling*.—Month of January, 1897. Estimated population, 36,000. Total deaths, 71; including phthisis pulmonalis, 9; enteric fever, 2; diphtheria, 2, and measles, 2.

Month of February, 1897. Total deaths, 68, including phthisis pulmonalis, 5; measles, 2, and whooping cough, 1.

Month of March, 1897. Total deaths, 66, including phthisis pulmonalis, 8; enteric fever, 3, and measles, 4.